

**COVID-19 (Coronavirus):
FAQs for Organ Transplantation**

Updated: August 9, 2021

The following FAQ was developed to relay information on the current state of knowledge. This document is subject to change as more information becomes available.

1. What is the origin of the novel coronavirus?

COVID-19 is the disease caused by the novel coronavirus named Severe Acute Respiratory Syndrome Coronavirus-2 (SARS-CoV-2) that was first recognized in the Hubei province of China in December 2019, subsequently spreading worldwide. COVID-19 was declared a pandemic on March 11, 2020. While the first infections with SARS-CoV-2 likely came from a non-human host, it quickly became well established in the human race by person-to-person transmission.

As of August 6, 2021, multiple [SARS-CoV-2 variants](#) as a result of viral mutation, are circulating globally. Variants of concern include B.1.1.7 (Alpha), B.1.351 (Beta), P.1 (Gamma), and B.1.617.2, (Delta) (1). These variants appear to be more transmissible (20-50% increased transmission) and some may reduce efficacy of vaccines and effectiveness of monoclonal antibodies.

2. How is SARS-CoV-2 transmitted?

There are 3 major ways the virus is spread: (1) inhalation of air carrying very small particles of infectious virus, (2) droplets of virus deposited onto exposed mucous membranes such as with being coughed on, (3) touching mucous membranes with hands that are contaminated with infected respiratory fluids.

Most frequently, transmission is presumed to be from via droplet spread from symptomatic individuals with COVID-19. However, shedding from asymptomatic and pre-symptomatic individuals can also transmit infection. The risk for exposure by breathing in the virus is felt to be highest when a person is within 3-6 feet of an infected individual. Aerosol generating events may also spread the infection. The incubation period is usually between 2-14 days in the general population although longer incubations have been documented (2).

Since asymptomatic individuals can spread the virus, the CDC recommends that unvaccinated people wear well fitting face masks that cover the nose and mouth when going out in public or

in instances where social distancing may be challenging. Given the data showing that vaccinated people can also spread the virus if infected, particularly with the delta variant, the CDC has now recommended that vaccinated people also wear masks if they are in an area with a high and substantial incidence of infection or if they are at high risk for complications from COVID-19 (3). Masks with exhalation valves or vents are not recommended as the hole in the material may allow escape of respiratory droplets(4).

Healthcare transmissions of COVID-19 have occurred. Given the potential for greater infectivity, strict isolation precautions should be followed for anyone with suspected SARS-CoV-2 infection. Healthcare personnel should all be vaccinated. In addition, regardless of vaccination status they should always wear a face mask while in the healthcare facility. Personal protective equipment should be used by all healthcare workers who enter the room of a patient with known or suspected COVID-19 or as specified by institutional policies. Surgical masks plus eye protection and an isolation gown are recommended. N95 masks or their equivalents should be used for procedures that are more likely to generate respiratory aerosolization (including bronchoscopy, intubation, and nasopharyngeal swab procurement). Local institutional guidelines should be followed for personal protective equipment (PPE).

3. Are transplant patients at higher risk for severe COVID-19?

Data on transplant recipients with COVID-19 suggests that although mild infections are common in transplant recipients, many have reported that infection, once acquired by immunosuppressed transplant recipients, may be of greater severity than in normal hosts. Experience with other viruses including prior outbreaks of coronaviruses, also suggests that severe infections can occur in some transplant recipients. The New York City experience revealed high rates of respiratory failure and mortality in transplant recipients but was likely impacted by the sudden and severe surge of infection that hit the city (5, 6). Data from other centers suggested similar outcomes in transplant recipients when compared to other higher risk populations when outcome analysis are corrected for comorbidities such as diabetes and hypertension, although this may vary based on organ transplant type (7-9).

It is important whenever a transplant patient has persistent fever or other symptoms of infection that they contact their transplant center for guidance. The CDC notes that the following adults of any age with the following conditions are at increased risk of severe COVID-19:

- Immunosuppressed after organ transplant
- Advanced age (over 60, but increasing with greater age)
- Hypertension
- Heart conditions, such as heart failure, coronary artery disease and cardiomyopathy
- Diabetes mellitus 2
- Obesity (BMI > 30)
- Cancer
- HIV infection

- Chronic kidney disease
- COPD
- Down syndrome
- Pregnancy
- Sickle cell disease
- Smoking

At this time, the risk factors for severe disease have been characterized by observational studies. An initial publication by Pereira, *et al* (6) reported advanced age was associated with severe disease in their transplant cohort, which is not different from reports of non-immunocompromised host. Kates, *et al* (10) reported age > 65, chronic lung disease, congestive heart failure, and obesity were independently associated with poor outcomes of COVID-19 in SOT. Further study is needed to better characterize which transplant recipients are at the greatest risk for poor outcomes.

4. Are there any specific travel restrictions for transplant patients?

The CDC recommends that travel be delayed unless fully vaccinated. However, vaccinated SOT recipients often do not mount as robust an immune response to vaccine when compared to the general population. We continue to recommend that transplant patients delay travel unless it is essential. This is both to avoid risks of transportation in non-private vehicles as well as risk of exposure in settings where COVID-19 rates might be higher. Should transplant recipients need to travel, we recommend taking additional essential medicines with them, to ensure they have a sustainable supply in the event of an unexpected quarantine or travel delay. For household members of SOT recipients, we recommend that they read the most up to date guidance on travel during COVID-19 at the CDC website (11, 12)

The CDC and WHO maintain websites that are being updated as the outbreak evolves, and travel recommendations will likely change over time.

- [CDC COVID Data Tracker](#)
- [World Health Organization COVID-19 Pandemic website](#)
- [Canada Public Health Website COVID-19](#)

5. Should transplant patients wear a mask or avoid public places?

The CDC has recently updated the guidance on masks for fully vaccinated individuals (3). Those with **immunocompromising conditions** are advised to continue safety precautions (i.e., mask wearing, frequent hand hygiene, and physical distancing), regardless of vaccinated status. They should avoid crowded places and reduce travel to areas of high SARS-CoV-2 prevalence as much as possible to reduce risk of exposure.

6. When should COVID-19 be considered in the differential diagnosis for transplant recipients?

Transplant patients with symptoms of a flu-like illness may have infection with SARS-CoV-2 as well as other infections. Many symptoms of COVID-19 are typical of respiratory viral infections. Transplant patients should be instructed to call the transplant center or their local physician if they have symptoms including, but not limited to, *fever, chills, rigors, cough, dyspnea, myalgias, headache, sore throat, diarrhea, or new loss of sense of taste and/or smell*. They should tell the transplant center if they have had close contact with a person known to have COVID-19 infection. They should notify the transplant center or hospital before presenting for care if possible. If patients are instructed to present for medical evaluation at a clinical center, transplant patients should wear a mask during transit and immediately upon entering the building. If the transplant patient has a medical emergency (e.g., shortness of breath, chest pain, or stroke/weakness), **they should call 911** and notify the dispatcher if they have been exposed to SARS-CoV-2 or have suggestive symptoms so that appropriate safety precautions can be taken.

There are many different causes for flu-like/respiratory symptoms. Although the prevalence of other respiratory viruses decreased dramatically with strict adherence to masking and social distancing, these illnesses are beginning to re-emerge with unmasking and decreased social distancing. Each hospital should have protocols in place for transplant patients with flu-like/respiratory symptoms in the era of COVID-19; these may vary seasonally in your geographic area. Consult your local hospital practices for outpatient COVID-19 screening availability or visitor restrictions for transplant recipients as these will change over time.

COVID-19 also has many [atypical clinical manifestations](#) affecting the skin (vesicles, rashes), cardiovascular system (e.g., myocarditis, cardiomyopathy, infarction), central nervous system (stroke and syncope, anosmia, dysgeusia), pulmonary emboli and vascular thrombosis, and renal or hepatic dysfunction (13). The clinician must consider testing for SARS-CoV-2 infection for individuals with these symptoms who are from areas where this infection is circulating.

Patients suspected of COVID-19 should wear a surgical mask, be placed in isolation and local infection control should be notified. CDC has updated the guidelines for infection control (14).

Specific testing for SARS-CoV-2 must be requested. Testing is done via a nucleic acid test or antigen detection assay (preferred test is RT-PCR on nasopharyngeal, nasal, mid-turbinate, or oropharyngeal swab) either as a single test or as part of a panel of tests for respiratory viruses. Testing guidelines vary by institution.

7. What is the approach to transplant candidates and recipients coming for routine appointments?

Each transplant program needs to decide their own policy for new transplants and outpatient visits when COVID-19 is circulating in the region.

Elective ambulatory appointments may be moved to virtual visits (e.g., telemedicine) and telephone contacts. Likewise, the urgency for bloodwork at the center, or for nonurgent procedures such as bronchoalveolar lavage and surveillance biopsies should be reviewed. Laboratory testing may be performed at centers outside the hospital or in the home if data can be provided expeditiously to the Transplant Center. Organizational leadership will need to be involved in prioritization plans. Some institutions may require SARS-CoV-2 testing prior to performing procedures on patients in both in and outpatient settings.

8. Should transplant candidates be counseled about potential risks for COVID-19 infection if called in for organ transplant?

Currently, it is appropriate to counsel all candidates about the risk for acquisition of COVID-19 from the community, the hospital environment and theoretically from an organ donor. Transplant candidates should be counseled on and encouraged to use preventive strategies (e.g., masking, physical distancing, frequent hand washing). Transplant candidates, professionals and household members should be encouraged to get vaccinated to reduce the risk of nosocomial transmission.

The risk-benefit ratio of transplantation during the COVID-19 pandemic should be reviewed with each patient considering individual risks of progression of underlying disease while on the wait-list and local infection and transmission rates. All deceased and living donors are tested for SARS-CoV-2 infection.

9. What is the approach to transplant candidates who are actively listed for transplant?

All patients in regions where SARS-CoV-2 is circulating should be tested for virus prior to transplantation, even if asymptomatic, regardless of vaccination status. It is not known if patients with active or recent COVID-19 can be safely transplanted. It is anticipated that transplantation of these patients with active viral infection and need for immunosuppression could result in adverse outcomes. The risk of transplantation must be balanced against the need for life-saving transplantation. Given limited treatment options for COVID-19, candidates with active COVID-19 should be deferred from transplantation, in most cases. Some patients continue to have positive PCR test for viral RNA long after symptoms have resolved (over 60 days from diagnosis in some patients). It is not clear if a persistently positive PCR test represents shedding of active virus or not. Blood tests that demonstrate antibodies to SARS-CoV-2 can be measured, but it is not yet known whether and how long the individual is protected against further infection or whether relapse could occur. There have been reports of reinfection in normal and immunosuppressed hosts.

The ideal disease-free interval before transplantation is unknown. Based on currently available data it is recommended that a candidate have complete symptom resolution and have a negative SARS-CoV-2 PCR from the respiratory tract prior to transplantation. This will also help to protect the hospital environment and the healthcare team. Some transplant physicians recommend two negative PCR tests at least 24 hours apart due to the limited sensitivity (~70%) of each test; the optimal timing of multiple tests is unknown. Likewise, the decision to proceed with transplant when disease-free but still with a positive test must be made on an individual basis weighing the likelihood of true active shedding versus death while awaiting another organ offer.

10. Are there any effective treatments for COVID-19?

For hospitalized patients with symptoms severe enough to require oxygen administration, a small number of therapeutic approaches appear to provide benefit. However, supportive care remains a vital component of patient care for both mild and severe disease.

Stable transplant patients with COVID-19 may be managed at home if they have social supports and access to medical care should the infection progress. In this setting, the main treatment is supportive care and consideration for monoclonal antibody administration as early as possible after symptom onset.

Telephone or video visits are an ideal approach for clinical monitoring. Home visits for laboratory testing or use of home oxygen saturation monitoring may be beneficial for outpatient assessment and determination of need for hospitalization.

A. Monoclonal Antibodies:

Current NIH and IDSA guidelines recommend that casirivimab/imdevimab or sotrovimab be used in ambulatory adults and pediatric patients with positive results of direct SARS-CoV-2 viral testing in the past 10 days and who are at high risk for progressing to severe COVID-19. The NIH no longer recommends use of bamlanivimab plus etesevimab as firstline therapy given concern for diminished effectiveness against potentially resistant viral variants(15).

Casirivimab and imdevimab should be administered together for the treatment of mild to moderate COVID-19 in ambulatory adults and pediatric patients. This combination was shown to reduce COVID-19-related hospitalization or emergency room visits in patients at high risk for disease progression within 28 days after treatment when compared to placebo (16). The casirivimab-imdevimab EUA was also recently expanded to use for post-exposure prophylaxis in immunocompromised or nonvaccinated patients (17, 18). Casirivimab and imdevimab are not authorized for patients who are hospitalized due to COVID-19 or require oxygen therapy due to COVID-19, and may be associated with worse clinical outcomes when administered to hospitalized patients with COVID-19 requiring high flow oxygen or mechanical ventilation. However, both combination therapies may be administered to patients who are hospitalized for reasons other than COVID-19 if they have evidence of mild disease and are at risk for progression to severe disease.

Sotrovimab is another monoclonal therapy now available by EUA for patients at risk for progression to severe COVID-19 (19). Authorization was based on interim analysis from a double-blind, placebo-controlled trial in which an 85% reduction in hospitalization or death was observed in patients who received sotrovimab compared with placebo. This monoclonal in addition is likely to provide protection against emerging viral variants, as it targets a highly conserved epitope in the receptor binding domain of SARS-CoV-2 spike protein.

B. Corticosteroids:

In patients requiring hospital admission with severe COVID-19 disease, the primary therapy with proven benefit in terms of mortality is corticosteroid therapy. In the RECOVERY trial, a controlled, open-label study from the UK, the incidence of death was lower in patients receiving dexamethasone (6mg daily for up to 10 days) compared with standard of care therapy in patients receiving mechanical ventilation (29.3% vs 41.4%), and to a lesser degree, in patients receiving oxygen therapy without mechanical ventilation (23.3% vs 26.2%) (20). The positive impact of corticosteroids on mortality in critically ill patients has been confirmed in a WHO meta-analysis as well as a meta-analysis of randomized-controlled trials (21, 22). Although specific information regarding inclusion or impact in transplant recipients has not been provided in these trials, this approach has now been used routinely in critically ill transplant recipients with COVID-19 where it appears to provide similar benefit as non-immunocompromised hosts. As hyperinflammation does occur in transplant recipients, the benefit of this therapy is not surprising; there remains concern, however, for promotion of secondary infections including bacterial and fungal pneumonias in immunocompromised patients.

C. Antiviral therapy:

While some trial data are encouraging, available antiviral agents appear to shorten the symptomatic period rather than eradicating viral infection. Physicians are encouraged to follow new or emerging data. Following our experience with antiviral therapy for influenza, start of antiviral therapy early in disease is more likely to be effective than administration later in the course of viral infection.

Remdesivir is an antiviral that inhibits viral genome replication and has been studied in a randomized controlled clinical trial for severe and moderate COVID-19 cases. The Adaptive COVID-19 Treatment Trial (ACTT)-1 Study revealed that remdesivir shortens the time to recovery in adults hospitalized with COVID-19 pneumonia (median 11 days vs 15 days), although the mortality rate did not significantly differ between groups (23). The FDA issued an Emergency Use Authorization (EUA) for remdesivir on May 1, 2020 to permit the emergency use of the unapproved product intravenously for treatment of suspected or laboratory confirmed COVID-19 in adults and children hospitalized with severe disease, defined as those with oxygen saturation (SpO₂) \leq 94% on room air or requiring supplemental oxygen or mechanical ventilation or extracorporeal membrane oxygenation. The drug was formally FDA approved October 22, 2020 (24). The SOLIDARITY multicenter trial of repurposed antiviral

agents also failed to show demonstrated impact on mortality in patients receiving remdesivir (25). This study, however, did not include immunocompromised patients, where the impact of antiviral therapy may be more likely to provide benefit in the setting of impaired immune response. Dosing is 200 mg IV on day one followed by 100 mg IV daily for a total of five days and liver function tests should be monitored during administration. While not recommended for use in patients with GFR<30 ml/min, emerging evidence suggests that remdesivir may be relatively safe in this group of patients (26). There is no transplant specific sub-analysis from this trial; data regarding the relative efficacy in SOT remain unavailable, although there does not seem to be increased toxicity in the setting of immunosuppressive drug use. In addition, additional case-control studies have demonstrated safety in terms of liver function elevations and acute kidney injury even in patients with impaired renal function (26).

D. Convalescent plasma:

A recent systematic review and meta-analysis of published and unpublished randomized controlled trials showed that there was no significant benefit of convalescent plasma on clinical outcomes (20), and analysis of data from the RECOVERY trial similarly failed to demonstrate benefit in terms of decrease in mortality (27). The current IDSA recommendation (28) is that this therapy be limited to patients participating in a clinical trial. High-titer antibody in convalescent plasma and early administration in course of illness may reduce hospitalization in cohort of outpatient older adults; hence, there may be a benefit in using convalescent plasma in patients with mild COVID-19 (29). Based on results of recent studies, the FDA updated the issued EUA for convalescent plasma, limiting the use of high-titer COVID-19 convalescent plasma for treatment of hospitalized patients early in the disease course (30). Thus far, there is no significant evidence of rejection or other safety signal related to the use of convalescent plasma(31, 32), although data remain limited in transplant recipients .

E. Tocilizumab:

Data on tocilizumab, an anti-IL-6 receptor monoclonal antibody, have been mixed, with initial randomized-controlled trials not meeting clinical endpoints of improved clinical status and mortality (33). However, analysis of two large multicenter trials, RECOVERY and REMAP-CAP, as well as additional randomized controlled trials, did demonstrate decreased mortality and lower risk of clinical deterioration when administered early in the hospital course in patients with rapidly progressive disease requiring mechanical ventilation or high flow nasal canula, in combination with dexamethasone (34-36). It remains unclear whether this treatment provides additional benefit, or poses additional risk, when administered to immunocompromised patients, especially in combination with dexamethasone, although case series describing its use do exist (37, 38). The NIH and IDSA guidelines recommend that tocilizumab should be used in patients with severe COVID who have elevated markers of systemic inflammation, in addition to standard of care that are not responding to dexamethasone or are within 72 hours of hospital admission and 24 hours of ICU admission, and are requiring ventilation or high flow nasal canula therapy. NIH guidelines recommend that tocilizumab administration be avoided “in patients who are significantly

immunosuppressed, particularly in those with recent use of other biologic immunomodulating drugs (15).”

F. Other approaches:

Prospective trials have been performed to examine the risks and benefits of other immunomodulatory therapies for the acute inflammatory state associated with severe COVID-19. As with corticosteroids and tocilizumab, concerns exist for use of such therapies when coupled with transplant immunosuppression due to the potential risk for superimposed infection. More data are required before a recommendation can be made. Reactivations of viruses such as hepatitis B virus (HBV), hepatitis C virus (HCV), or herpesviruses (HSV, CMV, VZV) or of latent infections such as tuberculosis or *Strongyloides* (39) may occur in patients receiving steroids or immunomodulation.

Several agents including interferon-1 β , leronlimab, kinase inhibitors, and anti-GMCSF antibodies were evaluated for anti-inflammatory responses; published data on efficacy have been variable. Sarilumab, another IL-6 inhibitor, failed to meet clinical trial endpoints, so there is insufficient data to recommend its use (40).

For all of these immune modulating therapies, few data are available from transplant recipients. Caution is warranted given the potential for increased vulnerability to secondary infections in patients receiving immunosuppression inhibiting the cellular immune response such as calcineurin inhibitors and mycophenolate mofetil. Given previous reports of allograft rejection with interferon therapy, we would recommend careful consideration of potential risks and benefits before enrollment of transplant recipients in interferon-based therapy trials.

A number of oral antiviral therapies are currently under investigation including the nitroxide TEMPOL, which inhibits the viral replicase, PF-07321332, a protease inhibitor, and molnupiravir, an RNA polymerase inhibitor. The availability of a safe and effective oral antiviral therapy that could be delivered early in infection in the outpatient setting would be extremely valuable for immunosuppressed patients at risk for progression to severe COVID-19 infection, but these therapies are currently in early stages of development and testing (19, 41).

Finally, NIH guidelines additionally recommend prophylactic dose anticoagulation for hospitalized patients, to prevent venous thromboembolism associated with COVID-19.

The following therapies are not recommended:

At this point in time, given negative data on the use of chloroquine and hydroxychloroquine, we do not recommend the use of these drugs, in accordance with National Institutes of Health (NIH) and IDSA guidelines (15, 28). The FDA EUA has been removed for chloroquine and hydroxychloroquine and these agents should be used with caution given known side effects. Patients receiving hydroxychloroquine require careful monitoring of QTc interval and for drug interactions.

Other therapies suggested for COVID-19 including colchicine (42) and ivermectin (43), have not shown significant clinical impact in adequately powered, randomized-controlled studies, and are not recommended at this time (15, 28).

G. Management of immunosuppression.

The impact of immunosuppression on COVID-19 is not currently known but decreasing immunosuppression may be considered for infected recipients who have not had recent rejection episodes. Many providers have decreased or discontinued cell cycle inhibitors or reduced calcineurin inhibitor levels, but comparative data regarding these interventions are not yet available. The decision to reduce immunosuppression should be based on severity of COVID-19 disease in comparison with rejection risk. Patients receiving maintenance corticosteroids have generally been maintained on these during therapy.

11. How do we approach clearance of transplant patients after COVID-19 infection for removal of enhanced isolation in the hospital and return to outpatient clinics?

Data suggests that in patients with mild to moderate COVID-19 infection, infectivity decreases within 10 days of symptom onset (44, 45), whereas recovery of viable virus is rare in patients with severe COVID-19 infection who are more than 20 days from symptom onset (46). Importantly, there is no evidence that adults with symptom resolution who have persistence of viral RNA transmit SARS-CoV-2 to their close contacts (47). Because of this, CDC no longer recommends a test-based strategy to determine when an individual with SARS-CoV-2 infection is no longer infectious.

It is unknown how long virus detected by sensitive PCR assays remains infectious in transplant recipients and what risk remains of infection for social contacts, healthcare providers, and the community in general.

In one study, 25% of kidney transplant recipients displayed persistent viral shedding more than 30 days after symptom onset (48). Additionally, PCR positivity has been reported in a heart transplant recipient 35 days after symptom onset, and in a kidney transplant recipient 63 days after symptom onset (49, 50). Viable virus, using cell culture, was identified in four kidney transplant recipients more than 3 weeks from symptom onset (8). Rare cases of prolonged infective virus have also been reported in a patient after CAR-T cell therapy for multiple myeloma (up to 119 days from infection onset) (51), a patient with lymphoma on B-cell directed therapy (52), as well as a patient with anti-phospholipid syndrome on rituximab and eculizumab with relapsing and recurring COVID-19 (53).

Patients *with prior positive SARS-CoV-2 PCR assays* should have resolution of symptoms, including fever, before enhanced respiratory precautions are removed. The CDC currently [recommends](#) that for mild or asymptomatic disease, patients must isolate for at least 10 days from date of symptom onset or diagnosis. In patients with severe infection or severe immunocompromise, isolation should be extended up to 20 days for “normal” hosts (54).

For organ transplant recipients with mild/asymptomatic COVID-19, it may be reasonable to follow CDC guidelines. However, some experts recommend adhering to a 20-day isolation with testing of those individuals at higher risk for prolonged infection, including those who have recently received cytolytic therapies or rituximab. The test-based strategy requires two negative PCR tests for SARS-CoV-2 from at least 2 consecutive respiratory samples collected more than 24 hours apart. It is unknown whether positive antibody testing (serology) is predictive of a protective immune response or of reduced infectivity, although this may be the case. Local institutional infection prevention policies should take precedence in removing patients from isolation.

If an elective procedure is required for a patient recovering from COVID-19, it may be advisable to wait until the patient is cleared and pre-procedural testing PCR testing is negative. For urgent procedures, strict infection control guidelines should be followed to minimize the risk of SARS-CoV-2 transmission to procedure teams. Ultimately, further data are required. Public health and local hospital guidelines should be followed.

12. Is a transplant recipient with resolved COVID-19 at risk for reinfection?

Cases of reinfection with COVID-19 have been reported but appear to be rare (55-57). A recent multicenter review reported a rate of reinfection of 0.2% (58). Whether reinfections result from deficient immune responses to the primary infection, are the result of waning immunity, or result from viral escape mutants remains to be determined. Hence, transplant recipients should continue to practice COVID-19 precautions (refer to #5) following natural infection with COVID-19 and after COVID-19 vaccination. It is especially important for close contacts (family, friends, co-workers) of transplant recipients to be vaccinated to avoid exposure to COVID-19.

13. Useful links:

Hygiene, physical distancing, masks, isolating, quarantining, what to do if you are sick:

- [CDC: How to Protect Yourself/What to Do If you are Sick](#)
- [CDC: Printable Handouts for Patients](#)

Treatment COVID-19 Guidelines

- [NIH Treatment Guidelines](#)

- [IDSA Treatment Guidelines](#)

Vaccines

- [American Society of Transplantation: COVID-19 Vaccine FAQ](#)

Drug-Drug interactions

- <https://www.covid19-druginteractions.org>

Donor Testing

- <https://www.myast.org/recommendations-and-guidance-organ-donor-testing>

REFERENCES

1. CDC. SARS-CoV-2 Variant Classifications and Definitions 2021 [updated July 20,2021. Available from: <https://www.cdc.gov/coronavirus/2019-ncov/variants/variant-info.html>.
2. Bai Y, Yao L, Wei T, Tian F, Jin DY, Chen L, et al. Presumed Asymptomatic Carrier Transmission of COVID-19. *Jama*. 2020;323(14):1406-7.
3. CDC. When You've Been Fully Vaccinated 2021 [updated May 16,2021. Available from: <https://www.cdc.gov/coronavirus/2019-ncov/vaccines/fully-vaccinated.html>.
4. CDC. Guidance for Wearing Masks (Guidance for Unvaccinated People) [updated April 19,2021. Available from: <https://www.cdc.gov/coronavirus/2019-ncov/prevent-getting-sick/cloth-face-cover-guidance.html>.
5. Akalin E, Azzi Y, Bartash R, Seethamraju H, Parides M, Hemmige V, et al. Covid-19 and Kidney Transplantation. *New England Journal of Medicine*. 2020;382(25):2475-7.
6. Pereira MR, Mohan S, Cohen DJ, Husain SA, Dube GK, Ratner LE, et al. COVID-19 in solid organ transplant recipients: Initial report from the US epicenter. *Am J Transplant*. 2020;20(7):1800-8.
7. Avery RK, Chiang TP, Marr KA, Brennan DC, Sait AS, Garibaldi BT, et al. Inpatient COVID-19 outcomes in solid organ transplant recipients compared to non-solid organ transplant patients: A retrospective cohort. *Am J Transplant*. 2020.
8. Chaudhry ZS, Williams JD, Vahia A, Fadel R, Parraga Acosta T, Prashar R, et al. Clinical characteristics and outcomes of COVID-19 in solid organ transplant recipients: A cohort study. *Am J Transplant*. 2020;20(11):3051-60.
9. Pereira MR, Arcasoy S, Farr MA, Mohan S, Emond JC, Tsapepas DS, et al. Outcomes of COVID-19 in solid organ transplant recipients: A matched cohort study. *Transpl Infect Dis*. 2021:e13637.
10. Kates OS, Haydel BM, Florman SS, Rana MM, Chaudhry ZS, Ramesh MS, et al. COVID-19 in solid organ transplant: A multi-center cohort study. *Clin Infect Dis*. 2020;2020 August 7:ciaa1097.
11. CDC. Domestic Travel During COVID-19 2021 [updated May 19,2021. Available from: <https://www.cdc.gov/coronavirus/2019-ncov/travelers/travel-during-covid19.html>.

12. CDC. International Travel During COVID-19 2021 [updated May 19,2021. Available from: <https://www.cdc.gov/coronavirus/2019-ncov/travelers/international-travel-during-covid19.html>.
13. Fishman JA, Roberts MB, Zhang EW, Kumar D, Hirsch HH, Maggiore U. Case 29-2020: A 66-Year-Old Man with Fever and Shortness of Breath after Liver Transplantation. *N Engl J Med*. 2020;383(12):1168-80.
14. CDC. Infection Control Guidance for Healthcare Professionals about Coronavirus (COVID-19) [updated June 3,2020. Available from: <https://www.cdc.gov/coronavirus/2019-ncov/infection-control.html>.
15. NIH COVID-19 Treatment Guidelines <https://www.covid19treatmentguidelines.nih.gov>: National Institutes of Health; [updated March 5,2021. Available from: <https://www.covid19treatmentguidelines.nih.gov/whats-new/>.
16. Weinreich DM, Sivapalasingam S, Norton T, Ali S, Gao H, Bhore R, et al. REGN-COV2, a Neutralizing Antibody Cocktail, in Outpatients with Covid-19. *N Engl J Med*. 2021;384(3):238-51.
17. FDA. FDA authorizes REGEN-COV monoclonal antibody therapy for post-exposure prophylaxis (prevention) for COVID-19 [Available from: <https://www.fda.gov/drugs/drug-safety-and-availability/fda-authorizes-regen-cov-monoclonal-antibody-therapy-post-exposure-prophylaxis-prevention-covid-19>.
18. O'Brien MP, Forleo-Neto E, Musser BJ, Isa F, Chan KC, Sarkar N, et al. Subcutaneous REGEN-COV Antibody Combination to Prevent Covid-19. *N Engl J Med*. 2021.
19. [Available from: <https://www.nih.gov/news-events/news-releases/nih-researchers-identify-potential-new-antiviral-drug-covid-19>.
20. Horby P, Lim WS, Emberson JR, Mafham M, Bell JL, Linsell L, et al. Dexamethasone in Hospitalized Patients with Covid-19 - Preliminary Report. *N Engl J Med*. 2020;Published online ahead of print.
21. Sterne JAC, Murthy S, Diaz JV, Slutsky AS, Villar J, Angus DC, et al. Association Between Administration of Systemic Corticosteroids and Mortality Among Critically Ill Patients With COVID-19: A Meta-analysis. *Jama*. 2020.
22. Pulakurthi YS, Pederson JM, Saravu K, Gupta N, Balasubramanian P, Kamrowski S, et al. Corticosteroid therapy for COVID-19: A systematic review and meta-analysis of randomized controlled trials. *Medicine (Baltimore)*. 2021;100(20):e25719.
23. Beigel JH, Tomashek KM, Dodd LE, Mehta AK, Zingman BS, Kalil AC, et al. Remdesivir for the Treatment of Covid-19 - Preliminary Report. *N Engl J Med*. 2020;E-pub ahead of print.
24. FDA. Fact Sheet for Health Care Providers Emergency Use Authorization of Veklury (remdesivir) 2020 [Available from: <https://www.fda.gov/media/137566/download>.
25. Consortium WHOIST, Pan H, Peto R, Henao-Restrepo AM, Preziosi MP, Sathiyamoorthy V, et al. Repurposed Antiviral Drugs for Covid-19 - Interim WHO Solidarity Trial Results. *N Engl J Med*. 2021;384(6):497-511.
26. Ackley TW, McManus D, Topal JE, Cicali B, Shah S. A Valid Warning or Clinical Lore: an Evaluation of Safety Outcomes of Remdesivir in Patients with Impaired Renal Function from a Multicenter Matched Cohort. *Antimicrob Agents Chemother*. 2021;65(2).
27. Horby PW, Estcourt L, Peto L, Emberson JR, Staplin N, Spata E, et al. 2021.
28. Bhimraj A, Morgan R, Hirsch Shumaker A, Lavergne V, Baden L, Chi-Chung Cheng V, et al. Infectious Diseases Society of America Guidelines on the Treatment and Management of Patients with COVID-19 <https://www.idsociety.org/practice-guideline/covid-19-guideline->

[treatment-and-management/](#): Infectious Diseases Society of America; [updated March 18,2021. Available from: <https://www.idsociety.org/practice-guideline/covid-19-guideline-treatment-and-management/>.

29. Libster R, Perez Marc G, Wappner D, Coviello S, Bianchi A, Braem V, et al. Early High-Titer Plasma Therapy to Prevent Severe Covid-19 in Older Adults. *N Engl J Med*. 2021.
30. FDA In Brief: FDA Updates Emergency Use Authorization for COVID-19 Convalescent Plasma to Reflect New Data [updated February 4,2021.
31. Rahman F, Liu STH, Taimur S, Jacobs S, Sullivan T, Dunn D, et al. Treatment with convalescent plasma in solid organ transplant recipients with COVID-19: Experience at large transplant center in New York City. *Clin Transplant*. 2020;34(12):e14089.
32. Fung M, Nambiar A, Pandey S, Aldrich JM, Teraoka J, Freise C, et al. Treatment of immunocompromised COVID-19 patients with convalescent plasma. *Transplant infectious disease : an official journal of the Transplantation Society*. 2021;23(2):e13477.
33. Roche provides an update on the phase III COVACTA trial of Actemra/RoActemra in hospitalised patients with severe COVID-19 associated pneumonia [updated July 29,2020. Available from: <https://www.roche.com/investors/updates/inv-update-2020-07-29.htm>.
34. Investigators R-C, Gordon AC, Mouncey PR, Al-Beidh F, Rowan KM, Nichol AD, et al. Interleukin-6 Receptor Antagonists in Critically Ill Patients with Covid-19. *N Engl J Med*. 2021;384(16):1491-502.
35. Group RC. Tocilizumab in patients admitted to hospital with COVID-19 (RECOVERY): a randomised, controlled, open-label, platform trial. *Lancet*. 2021;397(10285):1637-45.
36. Salama C, Han J, Yau L, Reiss WG, Kramer B, Neidhart JD, et al. Tocilizumab in Patients Hospitalized with Covid-19 Pneumonia. *N Engl J Med*. 2021;384(1):20-30.
37. Perez-Saez MJ, Blasco M, Redondo-Pachon D, Ventura-Aguiar P, Bada-Bosch T, Perez-Flores I, et al. Use of tocilizumab in kidney transplant recipients with COVID-19. *Am J Transplant*. 2020;20(11):3182-90.
38. Pereira MR, Aversa MM, Farr MA, Miko BA, Aaron JG, Mohan S, et al. Tocilizumab for severe COVID-19 in solid organ transplant recipients: a matched cohort study. *Am J Transplant*. 2020;20(11):3198-205.
39. Stauffer WM, Alpern JD, Walker PF. COVID-19 and Dexamethasone: A Potential Strategy to Avoid Steroid-Related Strongyloides Hyperinfection. *Jama*. 2020.
40. Sanofi provides update on Kevzara® (sarilumab) Phase 3 trial in severe and critically ill COVID-19 patients outside the U.S. [updated September 1,2020. Available from: <https://www.sanofi.com/en/media-room/press-releases/2020/2020-09-01-07-00-00>.
41. Pavan M, Bolcato G, Bassani D, Sturlese M, Moro S. Supervised Molecular Dynamics (SuMD) Insights into the mechanism of action of SARS-CoV-2 main protease inhibitor PF-07321332. *J Enzyme Inhib Med Chem*. 2021;36(1):1646-50.
42. Tardif J-C, Bouabdallaoui N, INVESTIGATORS C. Efficacy of Colchicine in Non-Hospitalized Patients with COVID-19. <https://www.medrxiv.org/content/101101/2021012621250494v1>. 2021.
43. Podder C, Chowdhury N, Sina M. Outcome of ivermectin treated mild to moderate COVID-19 cases: a single-centre, open-label, randomised controlled study. *IMC Journal of Medical Science*. 2020;14(2):002.
44. Wölfel R, Corman VM, Guggemos W, Seilmaier M, Zange S, Müller MA, et al. Virological assessment of hospitalized patients with COVID-2019. *Nature*. 2020;581(7809):465-9.

45. Rhee C, Kanjilal S, Baker M, Klompas M. Duration of Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2) Infectivity: When Is It Safe to Discontinue Isolation? *Clin Infect Dis*. 2021;72(8):1467-74.
46. van Kampen JJA, van de Vijver D, Fraaij PLA, Haagmans BL, Lamers MM, Okba N, et al. Duration and key determinants of infectious virus shedding in hospitalized patients with coronavirus disease-2019 (COVID-19). *Nat Commun*. 2021;12(1):267.
47. CDC. Interim Guidance on Ending Isolation and Precautions for Adults with COVID-19 2021 [updated March 16,2021. Available from: <https://www.cdc.gov/coronavirus/2019-ncov/hcp/duration-isolation.html>.
48. Benotmane I, Gautier-Vargas G, Wendling MJ, Perrin P, Velay A, Bassand X, et al. In-depth virological assessment of kidney transplant recipients with COVID-19. *Am J Transplant*. 2020;20(11):3162-72.
49. Man Z, Jing Z, Huibo S, Bin L, Fanjun Z. Viral shedding prolongation in a kidney transplant patient with COVID-19 pneumonia. *American Journal of Transplantation*. 2020.
50. Decker A, Welzel M, Laubner K, Grundmann S, Kochs G, Panning M, et al. Prolonged SARS-CoV-2 shedding and mild course of COVID-19 in a patient after recent heart transplantation. *American Journal of Transplantation*. 2020;n/a(n/a).
51. Hensley MK, Bain WG, Jacobs J, Nambulli S, Parikh U, Cillo A, et al. Intractable COVID-19 and Prolonged SARS-CoV-2 Replication in a CAR-T-cell Therapy Recipient: A Case Study. *Clin Infect Dis*. 2021.
52. Baang JH, Smith C, Mirabelli C, Valesano AL, Manthei DM, Bachman MA, et al. Prolonged Severe Acute Respiratory Syndrome Coronavirus 2 Replication in an Immunocompromised Patient. *J Infect Dis*. 2021;223(1):23-7.
53. Choi B, Choudhary MC, Regan J, Sparks JA, Padera RF, Qiu X, et al. Persistence and Evolution of SARS-CoV-2 in an Immunocompromised Host. *N Engl J Med*. 2020;383(23):2291-3.
54. CDC. Discontinuation of Transmission-Based Precautions and Disposition of Patients with COVID-19 in Healthcare Settings (Interim Guidance) [updated August 10,2020. Available from: <https://www.cdc.gov/coronavirus/2019-ncov/hcp/disposition-hospitalized-patients.html>.
55. Tomkins-Tinch CH, Daly JS, Gladden-Young A, Theodoropoulos NM, Madaio MP, Yu N, et al. SARS-CoV-2 Reinfection in a Liver Transplant Recipient. *Ann Intern Med*. 2021.
56. Klein J, Brito AF, Trubin P, Lu P, Wong P, Alpert T, et al. Case Study: Longitudinal immune profiling of a SARS-CoV-2 reinfection in a solid organ transplant recipient. *medRxiv*. 2021.
57. Babiker A, Marvil CE, Waggoner JJ, Collins MH, Piantadosi A. The Importance and Challenges of Identifying SARS-CoV-2 Reinfections. *Journal of clinical microbiology*. 2021;59(4).
58. Lawandi A, Warner S, Sun J, Demirkale CY, Danner RL, Klompas M, et al. Suspected SARS-CoV-2 Reinfections: Incidence, Predictors, and Healthcare Use among Patients at 238 U.S. Healthcare Facilities, June 1, 2020- February 28, 2021. *Clin Infect Dis*. 2021.